




Postpartum mood disorders

How to differentiate normal from problematic behaviors

	 Routine care Baby blues Offer reassurance	 Caution Mild to moderate depression Non-emergent evaluation	 Stop Severe depression or psychosis Prompt evaluation
Risk factors (see list on back)	Absent/mild	Present	Present
Self-reported symptoms	None or mild	Yes	Yes
Mood changes (sadness, anxious, irritable, empty feeling)	Sometimes	Persistent	Persistent
Feeling overwhelmed	Sometimes	Persistent	Persistent
Fatigue	Sometimes	Persistent	Persistent
Inability to sleep (even when baby sleeps)	No	Yes	Yes
Difficulty concentrating, remembering, making decisions	No	Yes	Yes
Feelings of guilt, worthlessness, "bad mom"	Sometimes	Persistent	Persistent
Lack of interest in self-care (bathing, dressing)	No	Yes	Yes
Difficulty bonding with baby	No/rarely	Yes	Yes
Worrying about baby	Sometimes	Persistent	Obsessively
Delusions	No	No	Yes
Hallucinations	No	No	Yes
Mania, paranoia	No	No	Yes
Feelings of harming self or baby or others	No	No	Yes
Being afraid of your thoughts	No	No	Yes
Use drugs or alcohol to relieve stress, elevate mood, help sleep, increase energy	Rarely	Often	Frequently/daily
Patient Health Questionnaire-2 (PHQ-2)			
Little interest or pleasure in doing things/withdraw from family/friends	None to several days	More than 50% of days	Nearly every day
Feeling down, depressed, hopeless	None to several days	More than 50% of days	Nearly every day
PHQ-9 score	0 to 5	6 to 15	> 15 or any score with a positive response to thoughts of suicide



Risk factors for postpartum mood disorders and substance use disorders:

- Mother is younger than 25
- Pregnancy was unplanned
- Mother has financial worries
- There is a lack of practical support (someone to help care for baby, run errands, take to doctor, etc.)
- Lack of emotional support (for example, from partner, family, or friends)
- History of birth trauma (real or perceived): bad outcome, unplanned outcome, NICU stay, negative emotions surrounding labor and delivery
- Difficult infant temperament
- History of postpartum depression with prior pregnancy

Helpful phone numbers:

- Postpartum Support International Help Line: **800-944-4773**
- National Maternal Mental Health Hotline: **833-852-6262** (call or text)
- National Suicide & Crisis Lifeline: **988**

Reference:

Postpartum Support International, <https://www.postpartum.net>

Anthem Blue Cross and Blue Shield is the trade name of Anthem Insurance Companies, Inc., independent licensee of the Blue Cross and Blue Shield Association. Anthem is a registered trademark of Anthem Insurance Companies, Inc. Providers who are contracted with Anthem Blue Cross and Blue Shield to serve Hoosier Healthwise, Healthy Indiana Plan and Hoosier Care Connect through an accountable care organization (ACO), participating medical group (PMG) or Independent Physician Association (IPA) are to follow guidelines and practices of the group. This includes but is not limited to authorization, covered benefits and services, and claims submittal. If you have questions, please contact your group administrator or your Anthem network representative.

INBCBS-CDCM-038013-23-CPN37940