

Maternal cardiovascular health

How to differentiate normal from problematic behaviors

| | ✓ | A | Ø |
|-----------------------------------|---|--|---|
| | Routine care Offer reassurance | Caution Non-emergent evaluation | Stop Prompt evaluation |
| Risk factors (see list on back) | Absent/mild | Present | Present |
| History of cardiovascular disease | None | None | Yes |
| Self-reported symptoms | None or mild | Yes | Yes |
| Shortness of breath | No interference with daily activities, with heavy exertion only | With moderate exertion, new onset asthma, persistent cough, or moderate to severe OSA | At rest, paroxysmal nocturnal dyspnew, bilateral chest infiltrates on CXR |
| Chest pain | Reflux related that resolves with treatment | Atypical | At rest or with minimal exertion |
| Palpitations | Few seconds, self-limited | Brief, self-limited episodes, no lightheadedness or syncope | Associated with near syncope |
| Syncope | Dizziness only with prolonged standing or dehydration | Vasovagal | Exertional or unprovoked |
| Fatigue | Mild | Mild to moderate | Extreme |
| Vital signs | Normal | Abnormal | Abnormal |
| Pulse (beats per minute) | < 90 | 90 to 119 | > 120 |
| Systolic BP | 120 to 139 | 140 to 159 | > 160 (or symptomatic BP) |
| Respirations (per minute) | 12 to 15 | 16 to 25 | > 25 |
| Oxygen saturation | > 97% | 95 to 97% | < 95% (unless chronic) |
| Physical exam | Normal | Abnormal | Abnormal |
| Jugular vein distention (JVD) | Not visible | Not visible | Visible > 2 cm above clavicle |
| Heart | S3, barely audible soft systolic murmur | S3, systolic murmur | Loud systolic murmur, diastolic murmur, S4 |
| Lungs | Clear | Clear | Wheezing, crackles, effusion |
| Edema | Mild | Moderate | Marked |



Risk factors for maternal cardiovascular disease:

- Mother is older than 35
- Mother is Black and non-Hispanic
- Mother has obesity
- Presence of a hypertensive disorder in pregnancy (preeclampsia, eclampsia, HELLP syndrome)
- Presence of chronic hypertension
- Presence of diabetes mellitus and gestational diabetes
- Mother has substance use disorder (in other words, uses alcohol, cocaine, methamphetamines)
- Mother uses tobacco
- · Strong family history of cardiovascular disease
- History of coronary artery dissection
- IUGR < 5%
- History of placental abruption
- Blood transfusion postpartum
- Peripartum infection

Medicaid services provided by Anthem Blue Cross, trade name of Blue Cross of California. Anthem Blue Cross and Blue Cross of California Partnership Plan, Inc. are independent licensees of the Blue Cross Association. Blue Cross of California is contracted with L.A. Care Health Plan to provide Medi-Cal Managed Care services in Los Angeles County. Commercial services provided by Anthem Blue Cross, trade name of Blue Cross of California. Anthem Blue Cross and Anthem Blue Cross Life and Health Insurance Company are independent licensees of the Blue Cross Association. Anthem is a registered trademark of Anthem Insurance Companies, Inc.